

Candidacy Exam  
Department of Physics  
January 20, 2007

Part I


Instructions:

- The following problems are intended to probe your understanding of basic physical principles. When answering each question, indicate the principles being applied and any approximations required to arrive at your solution. *If information you need is not given, you may define a variable or make a reasonable physical estimate, as appropriate.* Your solutions will be evaluated based on clarity of physical reasoning, clarity of presentation, and accuracy.
- Please use a new blue book for each question. Remember to write your name and the problem number of the cover of each book.
- We suggest you read all *four* of the problems before beginning to work them. You should reserve time to attempt every problem.

Fundamental constants:

Avogadro's number	$N_A$	$6.022 \times 10^{23} \text{ mol}^{-1}$
Boltzmann's constant	$k_B$	$1.381 \times 10^{-23} \text{ J K}^{-1}$
Electron charge magnitude	$e$	$1.602 \times 10^{-19} \text{ C}$
Gas constant	$R$	$8.314 \text{ J mol}^{-1} \text{ K}^{-1}$
Planck's constant	$h$	$6.626 \times 10^{-34} \text{ J s}$
	$\hbar = h/2\pi$	$1.055 \times 10^{-34} \text{ J s}$
Speed of light in vacuum	$c$	$2.998 \times 10^8 \text{ m s}^{-1}$
Permittivity constant	$\epsilon_0$	$8.854 \times 10^{-12} \text{ F m}^{-1}$
Permeability constant	$\mu_0$	$1.257 \times 10^{-6} \text{ N A}^{-2}$
Gravitational constant	$G$	$6.674 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$
Standard atmospheric pressure	1 atmosphere	$1.01 \times 10^5 \text{ N m}^{-2}$
Stefan-Boltzmann constant	$\sigma$	$5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$
Electron rest mass	$m_e$	$9.109 \times 10^{-31} \text{ kg} = 0.5110 \text{ MeV } c^{-2}$
Proton rest mass	$m_p$	$1.673 \times 10^{-27} \text{ kg} = 938.3 \text{ MeV } c^{-2}$
Origin of temperature scales	$0^\circ\text{C} = 273 \text{ K}$	

- I-1. A uniform ladder of mass  $m$  and length  $l$  leans against a frictionless vertical wall and rests on a frictionless horizontal floor. It is released from rest, with the ladder and the floor initially making an angle  $\theta_0$ .
- Derive the moment of inertia of the ladder about its center-of-mass in the plane in which the ladder is moving.
  - At some point, the ladder will separate from the *wall*. Determine the angle the ladder makes with the floor when this happens.



- I-2. A perfectly insulating hollow sphere of radius  $R$  has a charge  $Q$  uniformly spread on its surface. It spins with angular velocity  $\omega$ . Find the magnetic dipole moment.
- I-3. Derive the following equation<sup>1</sup> for the time-dependence of the expectation value of a (time-independent) quantum mechanical operator  $V$ :

$$\frac{d}{dt} \langle \psi | V | \psi \rangle = \frac{1}{i\hbar} \langle \psi | [V, H] | \psi \rangle, \quad (\text{I-1})$$

where  $H$  is the Hamiltonian.

Consider now an electron immersed in a uniform homogeneous field along the  $z$ -axis. This gives the following term in the Hamiltonian:

$$\frac{Be}{m} S_z, \quad (\text{I-2})$$


where  $B$  is the magnetic field,  $S_z$  is the  $z$ -component of the electron's spin operator,  $m$  is its mass, and  $e$  is the absolute value of the electron's charge. Assume that the remainder of the Hamiltonian is spin-independent.

---

<sup>1</sup>Note that there was a sign error in this equation on the exam as given; it has been corrected here.

Suppose that, at time  $t = 0$ , the electron is in a state with its spin component along the  $x$ -axis being  $\hbar/2$ . Determine the expectation values of the  $x$ ,  $y$  and  $z$  components of the electron's spin operators,  $\langle \psi | S_\alpha | \psi \rangle$ , at any later time. Qualitatively describe the time-dependence of the vector  $\langle \psi | \vec{S} | \psi \rangle$ .

- I-4. The diagram shows a spherical source of light illuminating a lens which focuses radiation onto a thin black heat-conducting disk. The light source is 1 mm in diameter and emits 100 W of radiation isotropically. The lens is 2 cm in diameter, has a focal length of 10 cm, and transmits all the radiation that impinges on it. The diameter of the disk is 0.5 mm and the image of the light source is at the disk and exactly covers the disk. Determine the equilibrium temperature of the disk. (See the table of constants for the value of the Stefan-Boltzmann constant.)



Candidacy Exam  
Department of Physics  
January 20, 2007

Part II


Instructions:

- The following problems are intended to probe your understanding of basic physical principles. When answering each question, indicate the principles being applied and any approximations required to arrive at your solution. *If information you need is not given, you may define a variable or make a reasonable physical estimate, as appropriate.* Your solutions will be evaluated based on clarity of physical reasoning, clarity of presentation, and accuracy.
- Please use a new blue book for each question. Remember to write your name and the problem number of the cover of each book.
- We suggest you read all *four* of the problems before beginning to work them. You should reserve time to attempt every problem.


Fundamental constants:

Avogadro's number	$N_A$	$6.022 \times 10^{23} \text{ mol}^{-1}$
Boltzmann's constant	$k_B$	$1.381 \times 10^{-23} \text{ J K}^{-1}$
Electron charge magnitude	$e$	$1.602 \times 10^{-19} \text{ C}$
Gas constant	$R$	$8.314 \text{ J mol}^{-1} \text{ K}^{-1}$
Planck's constant	$h$	$6.626 \times 10^{-34} \text{ J s}$
	$\hbar = h/2\pi$	$1.055 \times 10^{-34} \text{ J s}$
Speed of light in vacuum	$c$	$2.998 \times 10^8 \text{ m s}^{-1}$
Permittivity constant	$\epsilon_0$	$8.854 \times 10^{-12} \text{ F m}^{-1}$
Permeability constant	$\mu_0$	$1.257 \times 10^{-6} \text{ N A}^{-2}$
Gravitational constant	$G$	$6.674 \times 10^{-11} \text{ m}^3 \text{ kg}^{-1} \text{ s}^{-2}$
Standard atmospheric pressure	1 atmosphere	$1.01 \times 10^5 \text{ N m}^{-2}$
Stefan-Boltzmann constant	$\sigma$	$5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$
Electron rest mass	$m_e$	$9.109 \times 10^{-31} \text{ kg} = 0.5110 \text{ MeV } c^{-2}$
Proton rest mass	$m_p$	$1.673 \times 10^{-27} \text{ kg} = 938.3 \text{ MeV } c^{-2}$
Origin of temperature scales	$0^\circ\text{C} = 273 \text{ K}$	


- II-1. A azimuthally symmetric disk of radius  $R$  and mass  $M$  rolls without slipping on a horizontal surface. The momenta of inertia of the disk about its center is  $I$ , and a spring/springs are attached between the axis and a point B a distance  $d$  below the plane. The spring(s) have a length  $l_0$  when relaxed and the force exerted by the springs towards B is  $k(l - l_0)$ , where  $l$  is the distance between the center of the disk and the point of attraction. Find the frequency of small oscillations about the position of equilibrium. All motion is in a plane perpendicular to the horizontal surface.



- II-2. The region of space ( $x > 0, y > 0$ ) is bound on two sides by grounded conducting planes. A charge  $q$  is placed in this region at the position  $(x_0, y_0, 0)$ . Calculate the work required to move the charge from  $(x_0, y_0, 0)$  to  $(\infty, y_0, 0)$ .



- II-3. A particle of mass  $M$  is bound in a 1D infinite potential well of width  $L$  that is divided into two equal parts:



On the left side, the potential is zero, on the right side the potential is  $V_0$ . Everywhere else the potential is infinite. The initial wave function for the particle is given by the form

$$\psi(x) = \begin{cases} A_1 \sin(8\pi x/L) & \text{for } -\frac{L}{2} < x < 0, \\ A_2 \sin(4\pi x/L) & \text{for } 0 < x < \frac{L}{2}. \end{cases} \quad (\text{II-1})$$

- (a) Under what condition(s) on  $V_0$ ,  $M$ , and  $L$  is this an eigenfunction of energy for the 1D Schrödinger equation?
- (b) Determine values for  $A_1$  and  $A_2$  that give a normalized energy eigenfunction of the given form.
- (c) What is its energy?
- (d) Now assume that the two halves are suddenly separated to form two separate infinite wells, each of width  $L/2$ . (Imagine suddenly inserting an infinite delta-function potential barrier at the origin to separate the halves,  $V(x) \mapsto V(x) + \infty\delta(x)$ .) Determine the probabilities that the particle ends up in either separated well.

II-4. An isolated impurity site in a semiconductor is in thermal equilibrium with a reservoir at a chemical potential  $\mu$  and temperature  $T$ . It is modeled as follows: Up to two extra electrons can bind to the impurity. Each electron is the same spatial state (an  $s$ -wave orbital). The energy of a one electron state is  $E$  and the energy of a two-electron state is  $2E - V$ .

- (a) List the four possible electronic states in a standard basis of spin eigenstates.
- (b) Calculate the grand partition function for electrons at the impurity site and calculate the mean number  $\langle n \rangle$  of electrons.
- (c) For what value of  $\mu$  is  $\langle n \rangle = 1$ ?
- (d) An external magnetic field  $B$  is now applied, and interacts with electron spin in the usual way (“Zeeman term”). Calculate the spin susceptibility when  $\langle n \rangle = 1$ . Interpret the results when  $V \ll k_B T$  and when  $V \gg k_B T$ .